

-148-

What is claimed is:

add
1. A method for identifying a drug candidate as an HIV protease inhibitor potentially resistive against loss of inhibitory activity due to development of resistant strains of HIV, the method comprising the following steps:

5
Step A: determining whether the drug candidate has a binding activity with respect to HIV protease of less than 1 μ M;

10
Step B: determining whether the drug candidate has an inhibitory activity with respect to HIV protease of less than 1 μ M;

15
Step C: determining whether the drug candidate has a binding activity with respect to FIV protease of less than 1 μ M;

Step D: determining whether the drug candidate has an inhibitory activity with respect to FIV protease of less than 1 μ M; and then

20
Step E: if, in said Steps A, B, C, and C, the drug candidate is determined to have binding and inhibitory activities with respect to both HIV protease and FIV protease of less than 1 μ M, then selecting the drug candidate as the HIV
25
protease inhibitor potentially resistive against loss of inhibitory activity due to development of resistant strains of HIV.

000000-244000

-149-

2. A method for synthesizing a drug candidate for inhibiting HIV protease, the drug candidate including an N-terminus, a C-terminus, and an α -keto amide core structure linking the N-terminus and the C-terminus, the N-terminus including an aromatic amino acid residue selected from the group consisting of phenylalanine, tyrosine, and O-substituted tyrosine, the aromatic amino acid including a carbonyl group for linking to and incorporation into the α -keto amide core structure, the C-terminus including a heterocyclic ring having a ring nitrogen and one or more substitutions, the ring nitrogen of the C-terminus for linking to and incorporation into the α -keto amide core structure, the method comprising the following steps:

Step A: providing an N-terminus precursor identical to the N-terminus except that the carbonyl group is replaced by an α -hydroxyl acid group;

Step B: providing a C-terminus precursor identical to the C-terminus except that the ring nitrogen forms a secondary amine;

Step C: coupling the N-terminus precursor of said Step A to the C-terminus precursor of said Step B to form a drug candidate precursor identical to the drug candidate except that the α -keto amide core structure of the drug candidate is replaced by an α -hydroxyl amide core structure linking and incorporating the carbonyl group of the N-terminus and the ring nitrogen of the C-terminus; and then

Step D: oxidizing the α -hydroxyl amide core structure of the drug candidate precursor of said

0007712-070999

-150-

Step C for forming the α -keto amide core
structure and the drug candidate.

0907743-070999

-151-

3. A method for synthesizing a library of nxm drug candidates for inhibiting HIV protease, each of the nxm drug candidates including an N-terminus selected from n N-termini where n is two or greater, a C-terminus selected from m C-termini where m is two or greater, and an α -keto amide core structure linking the N-terminus and the C-terminus, each of the n N-termini including an aromatic amino acid residue selected from the group consisting of phenylalanine, tyrosine, and O-substituted tyrosine, the aromatic amino acid including a carbonyl group for linking to and incorporating into the α -keto amide core structure, each of the m C-termini including a heterocyclic ring having a ring nitrogen and one or more substitutions, the ring nitrogen of the C-terminus for linking to and incorporating into the α -keto amide core structure, the method comprising the following steps:

Step A: providing n N-terminus precursors

identical in structure to the n N-termini except that the carbonyl group of the N-termini is replaced by an α -hydroxyl acid group within the N-terminus precursors;

Step B: providing m C-terminus precursors

identical in structure to the m C-termini except that the ring nitrogen of the C-termini forms a secondary amine within the C-terminus precursors;

Step C: providing nxm reaction vessels;

Step D: loading each of the n N-terminus precursors into m of the reaction vessels of said Step C;

Step E: loading each of the m C-terminus

0007713-070990

-152-

precursors into n of reaction vessels of said
Step D for forming nxm admixtures of N-terminus
precursor and C-terminus precursors; then
Step F: within each of the nxm admixtures of said
Step E, coupling the N-terminus precursor to the
C-terminus precursor to form nxm drug candidate
precursors identical to the nxm drug candidates
except that the α -keto amide core structure of
the nxm drug candidates is replaced by an α -
hydroxyl amide core structure linking and
incorporating the carbonyl group of the N-
terminus and the ring nitrogen of the C-
terminus; and then
Step G: within each of the nxm reaction vessels,
oxidizing the α -hydroxyl amide core structure of
each of the nxm drug candidate precursors of
said Step F for forming the α -keto amide core
structure and the library of nxm drug
candidates.

4. A method for synthesizing a library of nxm drug
candidates for inhibiting HIV protease, each of the
nxm drug candidates including an N-terminus selected
from n N-termini where n is two or greater, a C-
terminus selected from m C-termini where m is two or
greater, and a hydroxyethylamine core structure
linking the N-terminus and the C-terminus, each of
the n N-termini including an aromatic amino acid
residue selected from the group consisting of
phenylalanine, tyrosine, and O-substituted tyrosine,
the aromatic amino acid including a hydroxyethyl
group in lieu of a carbonyl group for linking to and
incorporating into the hydroxyethylamine core

-153-

structure, each of the m C-termini including a heterocyclic ring having a ring nitrogen and one or more substitutions, the ring nitrogen of the C-terminus for linking to and incorporating into the hydroxyethylamine core structure, the method comprising the following steps:

Step A: providing n N-terminus precursors identical in structure to the n N-termini except that the hydroxyethyl group of the N-termini is replaced by an epoxide group within the N-terminus precursors;

Step B: providing m C-terminus precursors identical in structure to the m C-termini except that the ring nitrogen of the C-termini forms a secondary amine within the C-terminus precursors;

Step C: providing nxm reaction vessels;

Step D: loading each of the n N-terminus precursors into m of the reaction vessels of said Step C;

Step E: loading each of the m C-terminus precursors into n of reaction vessels of said Step D for forming nxm admixtures of N-terminus precursor and C-terminus precursors; then

Step F: within each of the nxm admixtures of said Step E, coupling the N-terminus precursor to the C-terminus precursor for forming the library of nxm drug candidates.

5. A library of nxm drug candidates for inhibiting HIV protease, each of the nxm drug candidates including an N-terminus selected from n N-termini

-154-

where n is two or greater, a C-terminus selected from m C-termini where m is two or greater, and an α -keto amide core structure linking the N-terminus and the C-terminus, each of the n N-termini including an aromatic amino acid residue selected from the group consisting of phenylalanine, tyrosine, and O-substituted tyrosine, the aromatic amino acid including a carbonyl group for linking to and incorporating into the α -keto amide core structure, each of the m C-termini including a heterocyclic ring having a ring nitrogen and one or more substitutions, the ring nitrogen of the C-terminus for linking to and incorporating into the α -keto amide core structure.

6. A library of nxm drug candidates for inhibiting HIV protease, each of the nxm drug candidates including an N-terminus selected from n N-termini where n is two or greater, a C-terminus selected from m C-termini where m is two or greater, and a hydroxyethylamine core structure linking the N-terminus and the C-terminus, each of the n N-termini including an aromatic amino acid residue selected from the group consisting of phenylalanine, tyrosine, and O-substituted tyrosine, the aromatic amino acid including a hydroxyethyl group *in lieu* of a carbonyl group for linking to and incorporating into the hydroxyethylamine core structure, each of the m C-termini including a heterocyclic ring having a ring nitrogen and one or more substitutions, the ring nitrogen of the C-terminus for linking to and incorporating into the hydroxyethylamine core structure.


-155-

7. An improved mechanism based inhibitor of HIV or FIV aspartyl protease of a type having an N-terminus, a C-terminus, and a core structure for linking the N-terminus to the C-terminus, the N-terminus including an aromatic amino acid residue linked to said core structure, the C-terminus including a heterocyclic ring including a ring nitrogen linked to said core structure, the core structure being isosteric with a scissile amide bond of a HIV or FIV aspartyl protease substrate, wherein the improvement comprises:
- 5 said core structure being an α -keto amide, and
- 10 the heterocyclic ring of said N-terminus being a pyrrolidine having at least one substituant other
- 15 than carboxylic acid and carboxymethyl ester.


0907712-070999

-156-

8. An improved mechanism based inhibitor of HIV or FIV aspartyl protease as described in claim 7 wherein said pyrrolidine is selected from the group represented by the following structures:




-157-




000713-0999

-158-


9. An improved mechanism based inhibitor of HIV or FIV aspartyl protease of a type having an N-terminus, a C-terminus, and a core structure for linking the N-terminus to the C-terminus, the N-terminus including an aromatic amino acid residue linked to said core structure, the C-terminus including a heterocyclic ring including a ring nitrogen linked to said core structure, the core structure being isosteric with a scissile amide bond of a HIV or FIV aspartyl protease substrate, wherein the improvement comprises:
- said core structure being an α -keto amide, and the heterocyclic ring of said N-terminus being a piperadine or an azasugar.
10. An improved mechanism based inhibitor of HIV or FIV aspartyl protease as described in claim 9 wherein said piperadine or azasugar is selected from the group represented by the following structures:




R = 85 H, 86 Me, 87 Bn




R = 88 H, 89 Me, 90 Bn



R = 91 H, 92 Me, 93 Bn



94




R = 95 H, 96 Me, 97 Bn

0907712-07009

11. An improved mechanism based inhibitor of HIV or
FIV aspartyl protease of a type having an N-terminus,
5 a C-terminus, and a core structure for linking the N-
terminus to the C-terminus, the N-terminus including
an aromatic amino acid residue linked to said core
structure, the C-terminus including a heterocyclic
ring including a ring nitrogen linked to said core
10 structure, the core structure being isosteric with a
scissile amide bond of a HIV or FIV aspartyl protease
substrate, wherein the improvement comprises:
said core structure being an α -keto amide, and
the aromatic amino acid of said C-terminus being
15 selected from a group consisting of tyrosine having a
protected amino, tyrosine having a protected amino
and a substituted hydroxyl, and phenylalanine having
a protected amino protected by carbobenzyloxy.
- 20 12. An improved mechanism based inhibitor of HIV or
FIV aspartyl protease as described in claim 11
selected from the group represented by the following
structures:

090774-0000

-160-



wherein R is selected from the group consisting of
hydrogen, hydroxy, benzyloxy, alkyl_(C1-C4)-oxy, o-
5 methoxy-benzyloxy, m-methoxy-benzyloxy, p-methoxy-
benzyloxy, o-methoxy-nitrobenzyloxy, m-methoxy-
nitrobenzyloxy, p-methoxy-nitrobenzyloxy, acetone,
benzylidene, 3-oxymethyl-catechol, 4-oxymethyl-
catechol; R₁ is selected from the group consisting of
10 carbobenzyloxy (CBZ), tert-butoxycarbonyl (t-BOC),
acyl; R₂ is selected from the group consisting of
hydrogen, benzyl, alkyl_(C1-C4), o-methoxy-benzyl, m-
methoxy-benzyl, p-methoxy-benzyl, o-methoxy-
nitrobenzyl, m-methoxy-nitrobenzyl, p-methoxy-
15 nitrobenzyl, 3-methylene-catechol, 4-methylene-
catechol.


09077712-070999

-161-

- 5 13. An improved mechanism based inhibitor of HIV or FIV aspartyl protease of a type having an N-terminus, a C-terminus, and a core structure for linking the N-terminus to the C-terminus, the N-terminus including an aromatic amino acid residue linked to said core structure, the C-terminus including a heterocyclic ring including a ring nitrogen linked to said core structure, the core structure being isosteric with a scissile amide bond of a HIV or FIV aspartyl protease substrate, wherein the improvement comprises:
- 10 said core structure being hydroxyethylamine, and the heterocyclic ring of said N-terminus being a pyrrolidine having at least one substituant other than carboxylic acid and carboxymethyl ester.
- 15
- 20 14. An improved mechanism based inhibitor of HIV or FIV aspartyl protease as described in claim 13 wherein said pyrrolidine is selected from the group represented by the following structures:

666020-2722050

-162-




0000202122060

15. An improved mechanism based inhibitor of HIV or FIV aspartyl protease of a type having an N-terminus, a C-terminus, and a core structure for linking the N-terminus to the C-terminus, the N-terminus including an aromatic amino acid residue linked to said core structure, the C-terminus including a heterocyclic ring including a ring nitrogen linked to said core structure, the core structure being isosteric with a scissile amide bond of a HIV or FIV aspartyl protease substrate, wherein the improvement comprises:

said core structure being hydroxyethylamine, and the heterocyclic ring of said N-terminus being a piperadine or an azasugar.





16. An improved mechanism based inhibitor of HIV or FIV aspartyl protease as described in claim 15 wherein said piperadine or azasugar is selected from the group represented by the following structures:



17. An improved mechanism based inhibitor of HIV or FIV aspartyl protease of a type having an N-terminus, a C-terminus, and a core structure for linking the N-terminus to the C-terminus, the N-terminus including an aromatic amino acid residue linked to said core structure, the C-terminus including a heterocyclic ring including a ring nitrogen linked to said core structure, the core structure being isosteric with a scissile amide bond of a HIV or FIV aspartyl protease substrate, wherein the improvement comprises:

said core structure being hydroxyethylamine, and the aromatic amino acid of said C-terminus being selected from a group consisting of tyrosine having a protected amino, tyrosine having a protected amino and a substituted hydroxyl, and phenylalanine having a protected amino protected by carbobenzyloxy.

18. An improved mechanism based inhibitor of HIV or FIV aspartyl protease as described in claim 17 selected from the group represented by the following structures:



09077712-03099

-165-

wherein R is selected from the group consisting of hydrogen, hydroxy, benzyloxy, alkyl_(C1-C4)-oxy, o-methoxy-benzyloxy, m-methoxy-benzyloxy, p-methoxy-benzyloxy, o-methoxy-nitrobenzyloxy, m-methoxy-nitrobenzyloxy, p-methoxy-nitrobenzyloxy, acetonide, benzylidene, 3-oxymethyl-catechol, 4-oxymethyl-catechol; R₁ is selected from the group consisting of carbobenzyloxy (CBZ), tert-butoxycarbonyl (t-BOC), acyl; R₂ is selected from the group consisting of hydrogen, benzyl, alkyl_(C1-C4), o-methoxy-benzyl, m-methoxy-benzyl, p-methoxy-benzyl, o-methoxy-nitrobenzyl, m-methoxy-nitrobenzyl, p-methoxy-nitrobenzyl, 3-methylene-catechol, 4-methylene-catechol.

0907712-070999